



Key

SADRU DAMJI
(Master)

**CHEMISTRY
HIGHER LEVEL
PAPER 1**

Tuesday 16 May 2000 (afternoon)

		1	2	3	4	5	6	7
1 hour	Grade							
	Boundaries	0-11	11-15	16-21	22-25	26-30	31-34	7, 35 out of 40

INSTRUCTIONS TO CANDIDATES

- Do not open this examination paper until instructed to do so.
- Answer all the questions.
- For each question, choose the answer you consider to be the best and indicate your choice on the answer sheet provided.

Periodic Table

Atomic Number																Atomic Mass	
1 H 1.01																	2 He 4.00
3 Li 6.94	4 Be 9.01															9 F 19.00	10 Ne 20.18
11 Na 22.99	12 Mg 24.31															17 Cl 35.45	18 Ar 39.95
19 K 39.10	20 Ca 40.08	21 Sc 44.96	22 Ti 47.90	23 V 50.94	24 Cr 52.00	25 Mn 54.94	26 Fe 55.85	27 Co 58.93	28 Ni 58.71	29 Cu 63.55	30 Zn 65.37	31 Ga 69.72	32 Ge 72.59	33 As 74.92	34 Se 78.96	35 Br 79.90	36 Kr 83.80
37 Rb 85.47	38 Sr 87.62	39 Y 88.91	40 Zr 91.22	41 Nb 92.91	42 Mo 95.94	43 Tc 98.91	44 Ru 101.07	45 Rh 102.91	46 Pd 106.42	47 Ag 107.87	48 Cd 112.40	49 In 114.82	50 Sn 118.69	51 Sb 121.75	52 Te 127.60	53 I 126.90	54 Xe 131.30
55 Cs 132.91	56 Ba 137.34	57 † La 138.91	72 Hf 178.49	73 Ta 180.95	74 W 183.85	75 Re 186.21	76 Os 190.21	77 Ir 192.22	78 Pt 195.09	79 Au 196.97	80 Hg 200.59	81 Tl 204.37	82 Pb 207.19	83 Bi 208.98	84 Po (210)	85 At (210)	86 Rn (222)
87 Fr (223)	88 Ra (226)	89 † Ac (227)	104 Rf (261)	105 Db (262)	106 Sg (263)	107 Bh (262)	108 Hs (262)	109 Mt (262)									


†

58 Ce 140.12	59 Pr 140.91	60 Nd 144.24	61 Pm 146.92	62 Sm 150.35	63 Eu 151.96	64 Gd 157.25	65 Tb 158.92	66 Dy 162.50	67 Ho 164.93	68 Er 167.26	69 Tm 168.93	70 Yb 173.04	71 Lu 174.97
--------------------	--------------------	--------------------	--------------------	--------------------	--------------------	--------------------	--------------------	--------------------	--------------------	--------------------	--------------------	--------------------	--------------------

†

90 Th 232.04	91 Pa 231.04	92 U 238.03	93 Np (237)	94 Pu (242)	95 Am (243)	96 Cm (247)	97 Bk (247)	98 Cf (251)	99 Es (254)	100 Fm (257)	101 Md (258)	102 No (259)	103 Lr (260)
--------------------	--------------------	-------------------	-------------------	-------------------	-------------------	-------------------	-------------------	-------------------	-------------------	--------------------	--------------------	--------------------	--------------------

1. According to the equation:



(volumes of gases)

what volume of air (20 % O_2) is required to react with 10 dm^3 of SO_2 ?

- A. 2 dm^3
 B. 5 dm^3
 C. 10 dm^3
 (D) 25 dm^3

$$\text{need } n_{\text{O}_2} = \frac{1}{2} n_{\text{SO}_2}$$

$$\Rightarrow \frac{1}{2} \times 10 \text{ dm}^3 = 5 \text{ dm}^3 \text{ O}_2$$

$$\begin{array}{lcl} 20 \text{ dm}^3 \text{ O}_2 & \text{in} & 100 \text{ dm}^3 \text{ AIR} \\ 5 \text{ dm}^3 \text{ O}_2 & \text{"} & 100 \times \frac{5}{20} = 25 \end{array}$$

2. Which of the following compounds has the greatest empirical formula mass?

		Emp. formula	E_r
A.	C_6H_6	CH	13
(B)	C_4H_{10}	C_2H_5	29
C.	C_3H_6	CH_2	14
D.	C_2H_6	CH_3	15


3. $\text{CaCO}_3(\text{s}) \rightarrow \text{CaO}(\text{s}) + \text{CO}_2(\text{g})$

When heated, CaCO_3 ($M_r = 100$) decomposes as shown above. When 20 g of impure CaCO_3 is heated, 0.15 moles of CO_2 are obtained. What is the percentage purity of the CaCO_3 ? (Assume that none of the impurities produce CO_2 upon heating.)

- A. 15
 B. 25
 C. 55
 (D) 75

$$\rightarrow 0.15 \text{ mol } \text{CaCO}_3 \text{ present} = 15 \text{ g}$$

$$\therefore \% = \frac{15}{20} \times 100 = 75\%$$




Chloroethene can be burned in oxygen as shown above. What is the value of w when $v = 2$?

A. 2

B. 3

C. 4

(D) 5



5. Which of the following particles contain more electrons than **neutrons**?

I. ${}^1_1\text{H}$ 1p 0n 1e⁻ ✓

II. ${}^{35}_{17}\text{Cl}^-$ 17p 18n 18e⁻ ✗

III. ${}^{39}_{19}\text{K}^+$ 19p 20n 18e⁻ ✗

(A) I only

B. II only

C. I and II only

D. II and III only

6. The first four ionisation energies (kJ mol^{-1}) for a particular element are 550, 1064, 4210 and 5500 respectively. This element should be placed in the same Group as

A. Li


(B) Be

C. B

D. C ← Chosen by many!

By jump
↑
2 valence e⁻
Group II/2

7. A certain element with two isotopes of masses M and $M+2$ is introduced into a mass spectrometer, vaporised and ionised. Which of the following paths are most likely for the resulting ions?



M $M+2$

A. I X IV

B. II X I X

C. IV III

D. IV II X

lighter therefore more deflected

8. A certain element has the electronic configuration $1s^2 2s^2 2p^6 3s^2 3p^6 4s^2 3d^1$. Which oxidation state(s) would this element most likely show?

A. +2 only

B. +3 only

C. +2 and +5 only

D. +2, +3, +4, +5

(should have been worded better)

← chosen by >50%

both 4s and 3d behave as valence electrons
4s² leave first, ∴ +2

1	1	1	1	1
---	---	---	---	---

then +3, +4, +5

9. Which one of the following increases in value from Li to Cs?

A. Atomic radius

B. Electronegativity

C. Ionisation energy

D. Melting point

down a group


decrease

10. Which of the following chlorides give neutral solutions when added to water?

I. NaCl ✓ SA + SB

II. Al_2Cl_6 ✗ Cl^- : neutral; Al^{3+} : weakly acidic ($\text{Al}(\text{H}_2\text{O})_6^{3+}$)

III. PCl_3 ✗ Hydrolyses to form H_3PO_3



(A) I only ✓

B. I and II only

C. II and III only ✗ *not the answer but chosen by many*

D. I, II and III

11. In which of the following is there at least one double bond?

I. O_2 $\ddot{\text{O}}=\ddot{\text{O}}$

II. CO_2 $\ddot{\text{O}}=\text{C}=\ddot{\text{O}}$

III. C_2H_4 $\begin{array}{c} \text{H} \quad \text{H} \\ \diagdown \quad \diagup \\ \text{C}=\text{C} \\ \diagup \quad \diagdown \\ \text{H} \quad \text{H} \end{array}$

A. I only

B. III only

C. II and III only

(D) I, II and III

12. According to VSEPR theory, which molecule would be expected to have the smallest bond angle?

(A) H_2O $\begin{array}{c} \ddot{\text{O}} \\ \diagup \quad \diagdown \\ \text{H} \quad \text{H} \end{array}$ $\approx 105^\circ$







B. H_2CO $\begin{array}{c} \text{H} \quad \text{H} \\ \diagdown \quad \diagup \\ \text{C}=\text{O} \end{array}$ 120°

C. CH_4 $\begin{array}{c} \text{H} \quad \text{H} \\ \diagdown \quad \diagup \\ \text{C} \\ \diagup \quad \diagdown \\ \text{H} \quad \text{H} \end{array}$ $109\frac{1}{2}^\circ$





D. NH_3 $\begin{array}{c} \text{H} \quad \text{H} \quad \text{H} \\ \diagdown \quad \diagup \quad \diagup \\ \text{N} \end{array}$ $\approx 107^\circ$

13. Which of the following can exist in both polar and non-polar forms?


poorly worded



14. What are the states of hybridisation for the carbon atoms in NCCH_2COOH ?



chosen by
is a third



15. Which of the following best accounts for the observation that gases are easily compressed?

A. Gas molecules have negligible attractive forces for one another.

☒ B. The volume occupied by the gas is much greater than that occupied by the molecules, almost $\checkmark$

C. The average energy of the molecules in a gas is proportional to the absolute temperature of the gas.

D. The collisions between gas molecules are elastic.

16. Which expression represents the density of a gas sample of relative molar mass, M_r , at temperature, T , and pressure, P ?

A. $\frac{PM_r}{T}$ (Unit do not match $\frac{1}{\text{mol}} \times M_r$)

$$PV = nRT$$

$$= \frac{m}{M} RT$$

B. $\frac{RT}{PM_r}$


$$\therefore \frac{PM}{RT} = \frac{m}{V} = d$$

☒ C. $\frac{PM_r}{RT}$

D. $\frac{RM_r}{PT}$

17.

Read question carefully



The heating curve for 10 g of a substance is given above. How much energy would be required to melt completely 20 g of the substance that is initially at 10° C?

→ Not ice ; (m.p.t = 30°C)

(A) 2400 J ↗ 20g



B. 1200 J

C. 800 J


D. 400 J

1200 J to melt 10g initially at 10°C
 $\therefore 2 \times 1200 \text{ J} \quad " \quad " \quad 2 \times 10 \text{ g} \quad " \quad " \quad "$

18.



Use the enthalpy values above to calculate ΔH for the reaction;






(A) -57 kJ





B. -114 kJ ✗

C. 57 kJ ✗





D. 114 kJ ✗



19. In which reaction is the change in entropy (ΔS) closest to zero?







20. The Born-Haber cycle for the formation of potassium chloride includes the steps below:







Which of these steps are exothermic?

poor
choices



21. Some collisions between reactant molecules do not form products. This is most likely because



22. Doubling which one of the following will double the rate of a first order reaction?

- (A) Concentration of the reactant
- B. Size of solid particles
- C. Volume of solution in which the reaction is carried out
- D. Activation energy

23. $F_2(g) + 2ClO_2(g) \rightarrow 2FClO_2(g)$

The following data were obtained for the reaction above. Use these data to determine the orders for the reactants F_2 and ClO_2 .

$[F_2(g)] / \text{mol dm}^{-3}$	$[ClO_2(g)] / \text{mol dm}^{-3}$	Rate / $\text{mol dm}^{-3} \text{s}^{-1}$
$\begin{matrix} 0.1 \\ \times 2 \rightarrow 0.2 \end{matrix}$	$\begin{matrix} 0.01 \\ \text{same} \rightarrow 0.04 \end{matrix} \times 4$	$\begin{matrix} 1.2 \times 10^{-3} \\ \times 4 \rightarrow 4.8 \times 10^{-3} \end{matrix} \therefore \propto [ClO_2]^1$
	$\begin{matrix} 0.01 \\ \rightarrow 0.01 \end{matrix}$	$\begin{matrix} 1.2 \times 10^{-3} \\ \times 2 \rightarrow 2.4 \times 10^{-3} \end{matrix} \therefore \propto [F_2]^1$

Order of reaction


	F_2	ClO_2
A.	1 ✓	1 ✓
B.	1 ✓	2 ✗
C.	2 ✗	1 ✓
D.	2 ✗	4 ✗


24. $2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g) \quad \Delta H = -197.8 \text{ kJ}$

The reaction above is an important step in the production of sulfuric acid. An increase in which of the following will increase the ratio of $\frac{SO_3(g)}{SO_2(g)}$ at equilibrium?

ie. favour the forward rxn

- A. Pressure only ✓
- B. Temperature only ✗ (decreased T will)
- C. Both temperature and pressure ✗ (chosen by some)
- D. Neither pressure nor temperature ✗







The equilibrium constant for the reaction above is 1.0×10^{-14} at 25°C and 2.1×10^{-14} at 35°C . What can be concluded from this information?

- A. $[\text{H}_3\text{O}^+]$ decreases as the temperature is raised. $\times$
- B. $[\text{H}_3\text{O}^+]$ is greater than $[\text{OH}^-]$ at 35°C . $\left. \begin{array}{l} \text{chosen by some} \\ \text{no} \end{array} \right\}$
- C. Water is a stronger electrolyte at 25°C .
- D.** The ionisation of water is endothermic. $\checkmark$

K increases with T
 $\therefore$ endothermic rxn



What is the equilibrium expression for the reaction above?

- A. $K_c = \frac{[\text{NH}_3]}{[\text{N}_2][\text{H}_2]}$ $\times$
- B. $K_c = \frac{2[\text{NH}_3]}{[\text{N}_2][\text{H}_2]}$ $\times$
- C. $K_c = \frac{2[\text{NH}_3]}{3[\text{N}_2][\text{H}_2]}$ $\times$
- D.** $K_c = \frac{[\text{NH}_3]^2}{[\text{N}_2][\text{H}_2]^3}$

$$K_c = \frac{[\text{NH}_3]^2}{[\text{N}_2][\text{H}_2]^3}$$

$\rightarrow$ too simple
 why not ask K_c' for the reverse rxn?

10 cm^3 of an HCl solution with a pH value of 2 was mixed with 90 cm^3 of water. What will be the pH of the resulting solution?

- A. 1
- B.** 3
- C. 5
- D. 7
- $\nwarrow$ chosen by $\approx 30\%$ of candidates


$$\therefore 10^{-2} \text{ mol dm}^{-3} \text{ HCl}$$

vol of sol goes from $10 \rightarrow 100\text{ cm}^3$; 10 fold

$\therefore$ conc decreases 10 fold (to 10^{-3})

$$\therefore \text{pH} = 3$$

ten fold dilution $\Rightarrow$ change in 1 pH unit



In the equilibrium above, what are the two conjugate bases?

- A. $\text{CH}_3\text{COOH}^{\times}$ and H_2O
- B. $\text{CH}_3\text{COO}^{\checkmark}$ and $\text{H}_3\text{O}^{+\times}$
- C. $\text{CH}_3\text{COOH}^{\times}$ and H_3O^+
- (D) $\text{CH}_3\text{COO}^{\checkmark}$ and $\text{H}_2\text{O}^{\checkmark}$

9. Which of the following is the weakest acid in aqueous solution?

- (A) $\text{C}_6\text{H}_5\text{OH}$ $K_a = 1.3 \times 10^{-10}$
- B. HCN $K_a = 4.9 \times 10^{-10}$
- C. H_2Se $K_a = 1.5 \times 10^{-4}$
- D. HF $K_a = 6.9 \times 10^{-4}$
- decreases
 } each choice
 chosen by about
 10% of
 candidates

10. Which salt will produce the most alkaline solution when dissolved in water?

- A. KNO_3
- B. MgCl_2
- (C) $\text{CH}_3\text{CO}_2\text{Na}$
- D. $(\text{NH}_4)_2\text{SO}_4$
- salts
 neutral from S.A. + S.B.
 $\text{CH}_3\text{COO}^- (\text{Na}^+) \leftarrow$ neutral
 conj. base of a weak acid
 NH_4^+ acidic
 SO_4^{2-} neutral anion of a S.A.

11. In the electrolysis of molten sodium chloride, the sodium ion goes to the

- A. positive electrode where it undergoes oxidation. $\times$
- B. negative electrode where it undergoes oxidation. $\times$
- C. positive electrode where it undergoes reduction. (chosen by some) $\checkmark$
- (D) negative electrode where it undergoes reduction. $\checkmark$
- Na^+ , cation to cathode; reduction

32. Which one of the following could reduce $\text{Cr}_2\text{O}_7^{2-}(\text{aq})$ to $\text{Cr}^{3+}(\text{aq})$?


*

- A. $\text{Ca}^{2+}(\text{aq})$ } highest ox. states
 B. $\text{Cu}^{2+}(\text{aq})$ }
 C. $\text{Fe}^{2+}(\text{aq})$ ← $\text{Fe}^{2+} \rightarrow \text{Fe}^{3+} + \text{e}^-$ need a reducing agent → it self oxidised
 D. $\text{Zn}^{2+}(\text{aq})$ highest ox. states

poorly done - only a 1/4 got this correct


33.

*



The standard electrode potentials for two metals are given above. What are the equation and cell potential for the spontaneous reaction that occurs?


- A. $\text{Tl}^+(\text{aq}) + \text{Cu}^{2+}(\text{aq}) \rightarrow \text{Tl}(\text{s}) + \text{Cu}(\text{s}) \quad E^\ominus = 0.003 \text{ V}$
 B. $2\text{Tl}(\text{s}) + \text{Cu}^{2+}(\text{aq}) \rightarrow 2\text{Tl}^+(\text{aq}) + \text{Cu}(\text{s}) \quad E^\ominus = 0.675 \text{ V}$
 C. $2\text{Tl}(\text{s}) + \text{Cu}^{2+}(\text{aq}) \rightarrow 2\text{Tl}^+(\text{aq}) + \text{Cu}(\text{s}) \quad E^\ominus = 1.011 \text{ V}$
 D. $2\text{Tl}^+(\text{aq}) + \text{Cu}(\text{s}) \rightarrow 2\text{Tl}(\text{s}) + \text{Cu}^{2+}(\text{aq}) \quad E^\ominus = 0.333 \text{ V}$



⇒ less than 1/2 got this correct

34. When molten magnesium chloride is electrolysed, how many moles of gaseous chlorine will be produced for every mole of magnesium?

A. $\frac{1}{2}$






B. 1 ✓

C. 2

D. 4



chosen by about 1/4 of the students

35. Which names are correct for the following isomers of C_6H_{14} ?



- A. I only
- B. I and II only
- C. I and III only
- D. I, II and III

Which of the compounds below will show ^{only one} ~~a~~ single peak in its $^1\text{H-NMR}$ spectrum?



(A) I only ✓


B. III only

C. I and II only

D. I, II and III

} chosen by $\approx 30\%$ of the candidates

What is the correct order of reaction types in the following sequence?



- | | I | II | III |
|-----|----------------|--------------|------------------|
| (A) | substitution ✓ | oxidation ✓ | esterification ✓ |
| B. | addition ✗ | substitution | substitution |
| C. | oxidation ✗ | substitution | addition |
| D. | substitution ✓ | oxidation ✓ | substitution ✗ |


Which carbon-containing product is most likely from the reaction of C_2H_4 and Br_2 ?

A. $\text{C}_2\text{H}_5\text{Br}$





(B) $\text{C}_2\text{H}_4\text{Br}_2$

C. $\text{C}_2\text{H}_3\text{Br}$






D. $\text{C}_2\text{H}_2\text{Br}_2$



39. Which of the following is expected to be a gas at 25° C?



40. Which of the compounds below is/are more likely to undergo substitution, rather than addition, reactions?



A. I only

B. II only

C. I and III only

D. II and III only ✓

typical rxn: Electrophilic aromatic substitution

chosen by about 1/3 of candidates